

continued

Within the next few weeks, representatives of the Women vs. Smoking Network appeared on *NBC Nightly News*, *CBS This Morning*, *CBS Evening News*, the *MacNeil-Lehrer NewsHour*, *Nightwatch*, and *Nightline*. Representatives were also interviewed by major national newspapers, including *USA Today*; by numerous local papers; by CBS Radio Network, the Black Radio Network, and National Public Radio; and by local talk shows. Last, representatives were asked to testify on the topic at congressional hearings. The network followed up on the publicity by spotlighting several different projects, including a petition to the tobacco companies to adhere to their own voluntary code of corporate ethics.

Even the cigarette's proposed name drew criticism. Groups in North Dakota and South Dakota objected to the name, as did Sioux tribal organizations, because "Dakota" means "friend" or "ally" in the Sioux language. These groups formed a coalition of more than 40 organizations and collected 25,000 signatures on a petition objecting to the use of the word and demanding that R.J. Reynolds cease

selling the cigarette, which had been test-marketed, as planned, beginning in April 1990. The Women vs. Smoking Network provided strategic counseling and technical support to the grassroots coalition and was instrumental in helping arrange a press conference in Washington, DC, in June 1990, which featured then Surgeon General Antonia Novello, Senator Larry Pressler (R-SD), and others objecting to the marketing plan.

Although advocacy groups were able to generate considerable community and media mobilization, R.J. Reynolds continued test-marketing. Advocates felt they had raised national concern about the targeting of cigarette advertising, although this impression was not directly verified through survey research. Dakota cigarettes were withdrawn two years later, however, because the brand did not sell as well as officials had hoped (*American Medical News* 1992). In this instance, although advocates might attribute the end result to the effective use of the media to promote the agenda for reducing tobacco use, the demise of the Dakota brand was probably more attributable to market forces.

chairman Joe Garagiola by lawmakers and Secretary of Health and Human Services Donna Shalala. More than \$70 million in electronic media coverage has been generated directly from NSTEP efforts. In addition, NSTEP activities appear to have substantially increased the coverage of smokeless tobacco issues in the print media. Before NSTEP there were approximately 500 print articles annually devoted to smokeless tobacco; since NSTEP that number has climbed to

more than 5,000. One article alone appeared in more than 800 newspapers on a given weekend, and NSTEP estimated the value of this media coverage at \$15 million. A recent survey of major league baseball players and coaches found that more than 44 percent of smokeless tobacco users want to quit in the next six months, perhaps attributable to NSTEP's active participation in educating ballplayers during spring training.

Components of Community Programs

Community Advocacy and Mobilization

Electronic Networking

Interactive communication technologies, such as computer networks, have been used extensively by advocacy groups for reducing tobacco use. For example, daily communications played an important

part in the response to Philip Morris' Bill of Rights Tour (see the text box later in this chapter). Many active, functioning networks now provide communication services to assist in efforts to reduce tobacco use.

The Institute for Global Communications, based in San Francisco, was an early provider of issue-specific networks to the general public. PeaceNet and

EcoNet, which were developed in 1986, are among the most widely used and well known of the institute's networks. As of October 1994, the institute reported a combined membership of 12,000 people from 130 countries (Moore 1994). Within these networks, and others like them, are smaller groups focused on a specific aspect of an issue or a particular policy. For instance, among HandsNet's 2,500 member organizations, which span the nonprofit sector, is a forum linking 200 community coalitions on substance abuse. This forum, managed by the Boston-based group Join Together and supported by the Robert Wood Johnson Foundation, provides on-line technical assistance to these coalitions. The forum also provides news summaries and information available on funding opportunities and proposed legislation.

Several networks link people who work in health-related areas. In 1993, the Public Health Network provided forums, e-mail service, and databases for its membership, which was composed of nearly 600 users from state and local health agencies and of program directors who were members of the CDC's Public Health Leadership Institute. In 1998, this network was replaced by the Information Network for Public Health Officials. Established by the CDC's Public Health Practice Program Office, the network links the public health community to the Internet and provides access to on-line information. Planned Parenthood Federation of America hosts PPXNet, a network for its affiliates in regional and national offices, primarily for communication within the organization itself. During the 1990s, the CDC offered the electronic resource WONDER to public health officials, academicians, and others so that they were able to communicate via e-mail with and have access to the CDC's databases of health data. The advent of the Internet, including Web-based e-mail and list serv technology, has facilitated the exchange of public health information for health professionals and the public. CDC now offers its health data, materials, databases, electronic journals, and other resources on its Web site at www.cdc.gov.

In 1990, the Advocacy Institute founded SCARCNet, a multiuser interactive bulletin board that served the tobacco control community. (The history of the bulletin board's sponsoring organization—the resource center known by the acronym SCARC—is discussed in “Impact of Direct Advocacy,” later in this chapter.) When SCARCNet ceased in January 2000, it had more than 1,000 subscribers and was circulated to thousands of readers throughout the world on various networks. SCARCNet's most popular feature was the “Daily Bulletin,” which each day summarized

major newspaper and journal stories on reducing tobacco use (Advocacy Institute 1994). The “Daily Bulletin” was accompanied by a “Morning Briefing,” which put these news stories in perspective for the tobacco control community. The contents of the “Daily Bulletin” stories were retained and stored in a database that is currently available for searching at www.tobacco.org. Another notable feature of SCARCNet was the publication of “Action Alerts.” These two-page summaries of current issues requiring immediate action included objectives for action, suggested actions, media bites, quotes, and talking points and were sent to SCARCNet as needed (on average, twice per month). The conferencing section on SCARCNet, called the “Strategy Exchanges,” provided a forum for planning, counseling, and experience sharing. The technology allowed for concurrent but separate discussions on discrete issues, such as clean indoor air, tobacco advertising and promotion, tobacco pricing policies, and minors' access to tobacco products. Since its inception in 1990 to its final edition on January 31, 2000, SCARCNet, along with its global counterpart GLOBALink, became an important resource for the tobacco control community. In February 2000, the American Legacy Foundation began its support of a newly designed and enhanced news service system that harnesses advances in Web technology to build on SCARCNet's valued features. This system provides users with the leading national news stories and also includes a news service that allows users to receive a customized selection of other stories based on their geographic location and specialty areas of greatest personal interest (e.g., advertising, enforcement, etc.).

SCARCNet has served as a model for other public health advocacy networks. Examples include Safety Net (an advocacy network for violence prevention) and the Marin Institute's ALCNet (a network for alcohol control advocates), which is modeled closely after SCARCNet. ALCNet has been used for media advocacy as well, particularly to facilitate strategy development to counteract certain alcohol products and promotions.

As with other modalities used for social change, the precise role of on-line networks—one element in a multifaceted approach—is difficult to define. Although process measures are available (e.g., frequency of interactions and message traffic), they do not assess the basic value of computer links in furthering the agenda for reducing tobacco use, nor is it likely (as is noted at the beginning of this chapter for social interventions overall) that their efficacy can be precisely estimated. Current enthusiasm for the mechanism,

however, will probably ensure its continuation, and accrued anecdotal experience—to date, quite positive—will provide the ultimate judgment.

Direct Advocacy

History and Activities

National-level activities, including the work of the Coalition on Smoking OR Health (see “Further Regulatory Steps” in Chapter 5; see also “Community Mobilization,” earlier in this chapter) and others (see Chapter 2 and USDHHS 1989b), have played a prominent role in the evolving policy changes concerning the reduction of tobacco use. Of equal interest, from the point of view of the potential impact of advocacy, are decentralized grassroots organizations.

The nonsmokers’ rights movement originated in the early 1970s (see “From Antismoking to Nonsmokers’ Rights” in Chapter 2). It consisted of individuals acting on their own and of small grassroots organizations of people irritated by ETS or convinced that their health suffered from it. During this period, the documented adverse health effects of ETS were first being brought to the public’s attention (Steinfeld 1972; U.S. Department of Health, Education, and Welfare 1972). As research documenting these health hazards accumulated, nonsmokers’ rights organizations grew in number and strength.

Many of the early grassroots organizations used the acronym GASP to represent similar titles, including the Group Against Smokers’ Pollution, the Group Against Smoking Pollution, the Group to Alleviate Smoking in Public Places, and Georgians Against Smoking Pollution. Other acronyms were also used, including FANS (Fresh Air for Nonsmokers), TAPS (Texans Against Public Smoking), and ANSR—pronounced “answer”—(Association for Nonsmokers Rights). Organizations were small, poorly funded, and often run from home by volunteers.

Initially, many nonsmokers’ rights organizations simply provided a forum for nonsmokers to express their concerns about smoking and ETS. These groups helped legitimize their members’ complaints and empower them to take protective actions. Such actions required courage, assertiveness, and no small measure of tact, since smoking in public areas was normative at the time. Group members might thus learn how to politely ask people to refrain from smoking; or to obviate direct confrontation with smokers, groups might provide members with signs, cards, or buttons asking people not to smoke in their presence.

Early in the movement, nonsmokers’ rights associations adopted public policy change as an

important goal. Groups began to work for passage of measures to restrict public smoking. Such regulations are often referred to as clean indoor air laws (see “Clean Indoor Air Regulation” in Chapter 5). To encourage these measures, an early GASP organization produced a “Bill of Rights” that stated, in part, that

Non-Smokers have the right to breathe clean air, free from harmful and irritating tobacco smoke. This right supersedes the right to smoke when the two conflict. Non-Smokers have the right to express—firmly but politely—their discomfort and adverse reactions to tobacco smoke. . . . Non-Smokers have the right to take action through legislative channels, social pressures or any other legitimate means—as individuals or in groups—to prevent or discourage smokers from polluting the atmosphere and to seek the restriction of smoking in public places (Group Against Smokers’ Pollution, n.d.).

Over time, many organizations moved to encompass broader policy goals for reducing tobacco use—in particular, they sought ways to decrease tobacco use by minors. Largely as a consequence of those efforts, direct advocacy and public policy change became important parts of these organizational strategies.

In some communities, nonsmokers’ rights organizations worked in isolation. In others, they formed associations with medical societies, voluntary health associations, and other organizations; the result was a more intense effort to ensure passage of desired legislation. Despite initial obstacles, in many communities nonsmokers’ rights associations were a driving force in moving their allies toward a legislative approach to reducing tobacco use. For example, one of the earliest and most influential nonsmokers’ rights organizations was California GASP, founded in 1976, which eventually became Americans for Nonsmokers’ Rights (ANR). ANR is now the principal national-level tobacco control group devoted primarily to promoting legislation for clean indoor air. In California, ANR helped support the passage of such ordinances in many localities. Partly as a result of ANR’s work, California has more local ordinances for clean indoor air than any other state. ANR has served as a national consultant to other groups pursuing such legislation.

Impact of Direct Advocacy

In retrospect, the grassroots organizations can be seen as having worked to diminish the legitimacy of tobacco use in the eyes of the public and the credibility of the tobacco industry. The passage of ordinances

against public smoking (see "Clean Indoor Air Regulation" in Chapter 5) occurred over several years, during which a shift in public opinion about smoking became evident. During the 1960s and 1970s, the right to smoke was largely unquestioned. In more recent years, declining smoking prevalence and public opinion polls have indicated an increasing intolerance for public smoking (USDHHS 1989b). The work of nonsmokers' rights organizations is coeval with these legal, epidemiologic, and social changes. Sorting out cause and effect is difficult, but the nonsmokers' rights movement seems to have contributed to the changing social norm (Glantz 1987).

There were, however, some important exceptions to the emerging nonsmoking norms. By the mid-1980s, it was apparent that both the traditional educational efforts and the passage of ordinances to protect nonsmokers from ETS had a limited effect on young people's smoking-related attitudes and behaviors (USDHHS 1994). Efforts to reduce smoking appeared unable to reduce the prevalence of smoking among teenagers (Lynch and Bonnie 1994), and smoking prevalence among white females began increasing sharply during the 1970s, as did the prevalence of smokeless tobacco use among males.

The failure to decrease smoking among young people is as difficult to assess as is the success observed among adults (particularly among adult men). Analyzing the effect of prevention activities on young people must include weighing the hampering effects of advertising and promotional efforts backed by the tobacco industry's enormous marketing budget (see "Advertising and Promotion" in Chapter 5; DiFranza et al. 1991; Pierce et al. 1991; Lynch and Bonnie 1994; USDHHS 1994). Whatever the interplay of the forces involved, the result is that protobacco activity directed at those entering the market has been generally successful. An exception is the continued decline in prevalence among young African Americans, particularly among young women (USDHHS 1998).

Perhaps some of the shortfall in grassroots efforts to reduce tobacco use is associated with the early isolation of these groups from the established national advocacy organization. Anecdotally, there is evidence of a culture clash. When the nonsmokers' rights movement emerged in the 1970s, many medical and voluntary health organizations decried what they perceived as the unprofessional, indecorous, confrontational approach that these activists took to an issue that had previously fallen in the domain of the traditional public health structure. Some traditional organizations in the public health arena may also have felt that

grassroots organizations were infringing on their "turf" and their fund-raising base.

For their part, nonsmokers' rights associations objected to what they saw as the overly cautious, measured approach of researchers, medical associations, and volunteer health associations, whose efforts seemed to have done little to solve the problems of day-to-day exposure to ETS. The grassroots organizations urged voluntary health organizations to examine their mission statements and dedicate appropriate resources to cost-effective solutions to reducing tobacco use.

In time, both approaches acknowledged that the lack of coordination and cohesion was a significant barrier to their efforts. The groups noted that, in contrast, the tobacco industry operated as a monolith through the coordinated efforts of the Tobacco Institute, a lobbying and public relations organization representing the industry. This insight led to the emergence of several groups—somewhat disparate in their approaches—that attempted to bridge some of the distance between the grassroots and national approaches to reducing tobacco use.

Among the oldest of these groups is DOC (Doctors Ought to Care), which was founded in 1977 as a national coalition of health professionals, students, and concerned individuals. DOC groups take an activist approach to public health problems and sponsor community projects and events on reducing tobacco use and other issues. From the outset, members chose confrontational programs, such as counteradvertising and picketing industry-sponsored sports events, to delegitimize the tobacco industry and focus attention on its activities by involving both physicians and young people in advocacy activities. DOC groups use satire, ridicule, and parody in their work to appeal to children and teenagers (Blum 1982); for example, they have sponsored "Emphysema Slims" tennis matches featuring appearances by "Martina Nosmokanova." DOC also maintains a large archive of activities related to the tobacco industry, including past advertising campaigns and marketing strategies (Mintz 1995). The activities of DOC are similar in style, if not content, to those of the Australian organization Billboard Utilising Graffitiists Against Unhealthy Promotions (BUGA-UP), which was founded in 1979. BUGA-UP members, some of whom are physicians, have used unconventional tactics, such as spray-painting billboards that advertise tobacco products (Jacobson 1983).

Another group is Stop Teenage Addiction to Tobacco (STAT), which was founded in 1985 with the aim of reducing tobacco use among minors. From its inception, STAT aimed to unite the medical and

scientific arm and the grassroots arm of the movement to reduce tobacco use. Although STAT frequently approaches tobacco issues from the activist perspective, the organization has long included key members of the medical and public health establishment in its leadership. DOC, STAT, and other groups have attempted to make the activist, confrontational approach to reducing tobacco use acceptable to the more conservative medical and voluntary health organizations. Partly because of these efforts, an activist approach is now an important component of the movement (see the text box "Bill of Rights Tour").

Another impetus for a more unified movement was the establishment of the Smoking Control Advocacy Resource Center (SCARC) at the Advocacy Institute in 1987. The Advocacy Institute's mission—to study, analyze, and teach public interest advocacy—included a focus on smoking reduction as a model public interest movement. The institute received funding from the Henry J. Kaiser Family Foundation to establish SCARC. Rather than be a frontline organization, SCARC proposed to help build the movement's infrastructure. As such, SCARC would be viewed as a neutral player and would not vie with the movement's other organizations in seeking media, voluntary, or funding sources. Since its formation, SCARC has served three important roles as convener, tobacco industry monitor, and center for strategic development, training, and counseling (Butler 1990).

Media Advocacy

Media advocacy for reducing tobacco use was developed during the 1980s by a small number of activists working primarily in the United States, Canada, Australia, and the United Kingdom. The attendees at the September 1985 International Summit of Smoking Control Leaders resolved to produce a handbook that would provide guidance on using the media to support tobacco control. The resulting document, *Smoke Signals: The Smoking Control Media Handbook* (Pertschuk 1987), describes many of the important themes and skills needed for using what would later be dubbed "media advocacy." In January 1988, the Advocacy Institute convened a two-day consensus workshop, sponsored by the NCI, that produced a second handbook on media advocacy, *Media Strategies for Smoking Control: Guidelines* (USDHHS 1989a), which formally recognized the importance of media advocacy in reducing tobacco use (and in which the term "media advocacy" was first employed).

Media advocacy has been defined as the strategic use of mass media to advance a social or public

policy initiative (NCI 1991). In contrast to the goal of traditional health communications efforts, the goal of media advocacy is to change public policy and thereby generate a broader impact on tobacco use by creating an environment in which smoking is not normative. *Smoke Signals* articulates six critical tasks the media must perform to help accomplish this goal: (1) educate the public about the severity of the risks of smoking, the susceptibility of every smoker, and the health benefits of quitting; (2) educate the public about the health risks of ETS; (3) alert citizens and policymakers to injurious public policies that promote smoking, including insufficiently regulated advertising and promotion of cigarettes, as well as unrestricted smoking in public areas and the workplace; (4) respond to and counteract the propaganda and disinformation campaigns of the tobacco industry; (5) counter the economic and political influence of the tobacco industry, which thwarts the adoption of remedial policies; and (6) reinforce evolving social nonsmoking norms (Pertschuk 1987).

Media advocacy campaigns have been likened to political campaigns "in which competing forces continuously react to unexpected events, breaking news, and opportunities" (Pertschuk et al. 1991, p. 3). Such campaigns require both presenting the public health side of an issue and negating the opposing side. Like political campaigns, media advocacy campaigns require quick reactions that contrast with the carefully planned, fixed agendas of traditional media programs.

Media advocacy recognizes the potential of the press to place on the public agenda issues concerning the reduction of tobacco use and to either advance or retard progress toward policy goals. Successful media influence requires gaining access to the news and framing or shaping coverage of the resulting story. These strategies are interrelated, since the framing of a story helps determine whether a journalist will agree to cover it.

The use of media advocacy has two daunting limitations: it is a new technique that requires complex skills and an understanding of the news media, and it demands a large investment in time (Wallack 1990). But another apparent barrier—the reliance on an outside party (the media) to achieve program goals—is also a source of considerable strength: media advocacy is a means by which public health practitioners can indirectly confront and compete with forces that are traditionally beyond their policy and financial reach. These forces represent powerful vested interests—the tobacco industry, advertising industry, retail establishments that sell tobacco, and others. The financial and political influence of these entities can limit the ability of public

Bill of Rights Tour

In fall 1989, Philip Morris, the largest U.S. manufacturer of cigarettes, contracted with the U.S. National Archives and Records Administration to sponsor a commemoration of the 200th anniversary of the Bill of Rights. The commemoration involved a national advertising campaign, including commercials on prime-time television and full-page advertisements in major newspapers, asking Americans to "Join Philip Morris and the National Archives in celebrating the 200th anniversary of the Bill of Rights" (cited in Advocacy Institute 1989, p. 1). Philip Morris soon announced plans to transport Virginia's copy of the Bill of Rights to all 50 states in cooperation with the Virginia State Library and Archives.

Advocates for reducing tobacco use interpreted Philip Morris' effort as an attempt to link smoking with the national freedoms guaranteed by the Bill of Rights. These groups believed that Philip Morris would use its association with the Bill of Rights Tour, which highlighted themes of liberty and freedom of expression, to gain public support for the company's claim of a First Amendment right to advertise. Philip Morris' project with the National Archives raised concern in the U.S. House of Representatives, which held hearings on the issue but did not intervene. Advocates for reducing tobacco use began using the 16-month tour schedule to coordinate local efforts to counter what they considered to be a tobacco-marketing plan.

The Washington state chapter of Doctors Ought to Care (DOC) built a countersymbol, the "Statue of Nicotina," to travel with the tour. At a press conference, comments from the president of the chapter, Dr. Robert Jaffe, captured the flavor of the symbol's proposed use:

Nicotina is modeled [on] the Statue of Liberty. She's holding a cigarette in her upheld hand, instead of a torch, and her eyes are closed, the symbol of shame that she's been . . . made a symbol of tobacco. The chains from her cigarettes in the pack help to illustrate to all of the children who are going to see the Bill of Rights Tour that this is a dangerous, addictive drug. At her feet are the words, "Give me your poor, your tired, your women, your children yearning to breathe free . . ." (quoted in Wallack et al. 1993, p. 185).

The Advocacy Institute published an advance schedule of the national tour, including dates and specific locations for each of the tour's stops. The institute also tracked activities in various states and disseminated strategic information through *Action Alerts* posted on SCARCNet, the institute's computer network dedicated to sharing information on reducing tobacco use. SCARCNet (see "Electronic Networking," earlier in this chapter) was a key mechanism for advocates to share information and develop strategies. In addition, the American Lung Association and the American Medical Association provided materials and strategic support to its interested affiliates.

Initially, Philip Morris responded to protests at tour sites by establishing a "speaker's corner" that restricted protesters to a site away from the exhibit hall. At first, this strategy successfully muted attacks and deflected positive attention from protesters. Indeed, by appearing to encourage protesters, Philip Morris was portrayed by some media reports as being faithful to the spirit of the Bill of Rights. As the tour continued, however, groups opposed to the sponsorship learned from experience in other states. The groups refined their message, learned how best to respond to Philip Morris' spokespersons, discussed public reaction to their protests, and modified their tactics appropriately. They developed a simple slogan, "Bill of Rights Yes/Philip Morris No" (cited in Wallack et al. 1993, p. 186), to clarify the theme of their protests.

With the changed approach, advocates reported improved media coverage of the protests. At almost every tour stop, advocates staged press conferences before the opening of the exhibit and displayed the Statue of Nicotina, which was transported from state to state. By February 1991, five months into the tour, Philip Morris scaled down the number of scheduled stops. The tour, accompanied by advocates for reducing tobacco use, continued through its conclusion in Richmond, Virginia, in December 1991.

The ultimate effectiveness of this advocacy effort is difficult to judge, but the effort played an obvious role in muting the public relations benefits to the tobacco industry. At the very least, the resources invested by the industry did not appear to bring the expected return.

(as well as private) agencies to use confrontational tactics. In addition, many communities prefer consensus building to confrontation with powerful opposition parties. However, because the visible products of media advocacy—the media reports themselves—emerge from a disinterested party (the media) rather than from parties for or against reducing tobacco use, this newest form of social intervention can be successful in previously problematic areas.

As with other social interventions, the precise contribution of media advocacy to the effort to reduce tobacco use is difficult to judge. Events like those surrounding the marketing of the cigarette brands Uptown, X, and Dakota and the Philip Morris-sponsored Bill of Rights Tour demonstrate the role that media advocacy can play in the overall effort.

Countermarketing

Mass Media in Tobacco Control

In contemporary society, the mass media are the most important means of educating and informing the public and, through public response to media, policymakers. By design or not, the media plays an enormous role in influencing the smoking behavior of individuals and the actions of policymakers in both the public and the private sector (Pertschuk 1987). Public health programs have used various health communication programs to inform and influence the behavior of the general public. Traditionally, communication programs intended to reduce tobacco use have tried to influence the behavior of individuals. Most such media campaigns have focused on influencing the behavior of adult smokers—and hence have focused more on smoking cessation than on prevention. Flay (1987) describes three prominent types of mass media programs and campaigns designed to influence smoking-related knowledge, attitudes, and behavior: (1) those that inform the public of the negative health consequences of cigarette smoking and try to motivate smokers to quit, (2) those that promote specific smoking cessation actions to those smokers motivated to quit (e.g., smokers are encouraged to call a help line or to request specific materials, such as a tip sheet or a self-help manual), and (3) those that promote smoking cessation self-help clinics for those smokers who desire to quit. A smaller number of campaigns have focused on youth, either encouraging young people to avoid using tobacco products or convincing young people who smoke to try to quit (USDHHS 1994).

A factor that has limited the success of traditional mass media campaigns is the small size of the campaign budgets compared with the advertising and

marketing budgets of the tobacco industry (Flay 1987; USDHHS 1994). In addition, these campaigns to reduce tobacco use have experienced drawbacks because of their traditional reliance on public service announcements (PSAs). Although PSAs have been an integral part of such efforts for many years, the number of PSAs on any subject provided to broadcasters has increased, whereas the amount of donated air time available for PSAs has decreased. Also, the advent of cable technology, which has increased the number of channels through which people can be reached and therefore has diffused the audience, has further hampered efforts to reach targeted groups efficiently. By the mid-1980s, it had become apparent that the role of the media in the effort to reduce tobacco use required reevaluation. In the following sections, the uses of mass media approaches for tobacco control are summarized.

Effects of Protobacco Advertising and Promotion

The effect of tobacco advertising and promotion activities on both adult consumption and youth initiation has been the subject of considerable research over the past decade (see “Advertising and Promotion” in Chapter 5). While noting that existing evidence suggests that tobacco marketing increases the level of tobacco consumption, the 1989 Surgeon General’s report *Reducing the Health Consequences of Smoking: 25 Years of Progress* concluded that the issue is so complex that a sufficiently rigorous study capable of providing definitive scientific evidence is not available and that “none is likely to be forthcoming in the foreseeable future” (USDHHS 1989b, pp. 516–7). The 1994 Surgeon General’s report *Preventing Tobacco Use Among Young People* similarly noted the absence of a definitive longitudinal study of the direct relationship of tobacco advertising to adolescent smoking. However, acknowledging the value of recent nonlongitudinal studies focused on young people, the report offered this major conclusion: “Cigarette advertising appears to increase young people’s risk of smoking by affecting their perceptions of the pervasiveness, image, and function of smoking” (USDHHS 1994, p. 6). Also in 1994, the Institute of Medicine concluded that the preponderance of evidence suggests that tobacco marketing encourages young people to smoke (Lynch and Bonnie 1994).

In its rule to restrict the access and appeal of tobacco products to young people, the Food and Drug Administration (FDA) reviewed the quantitative and qualitative evidence and concluded that cigarette advertising is causally related to the prevalence of smoking among young people (*Federal Register* 1996). The

agency also cited statements from internal documents of the tobacco industry to show the importance of the youth market segment to the industry's continued success. More recently, a 1998 Report to the United Kingdom's Chief Medical Officer by the Scientific Committee on Tobacco and Health concluded unanimously that tobacco advertising and promotion influence young people to begin smoking (Scientific Committee on Tobacco and Health 1998).

Survey data show that among children who smoke, most use the most heavily advertised brands of cigarettes, whereas many adult smokers buy generic or value category brands, which have little or no image advertising (CDC 1994). A major econometric marketing study found that young people are three times more affected by advertising than are adults (Pollay et al. 1996). Research has also pointed to the impact of other tobacco promotional activities, such as sponsorship of public entertainment events and distribution of specialty or premium items. These activities constitute the largest (and an increasing) share of tobacco marketing expenditures. The CDC has estimated that today's U.S. teens already have been exposed to more than \$20 billion in imagery advertising and promotions since age 6, creating a "friendly familiarity" for tobacco products and an environment in which smoking is seen as glamorous, social, and normal (Eriksen 1997). Although the effect of this exposure is difficult to quantify, especially nationwide, one study has estimated that 34 percent of all youth experimentation with smoking in California between 1993 and 1996 can be attributed to tobacco promotional activities (Pierce et al. 1998). A recent study found that teenagers who can readily name a cigarette brand and who own a tobacco-company-sponsored promotional item are more than twice as likely to become established smokers than adolescents who do neither (Biener and Siegel 2000).

Effects of Tobacco Countermarketing

In light of ubiquitous and sustained protobacco messages, countermarketing efforts of comparable intensity and duration are needed to alter the social and environmental context of tobacco use. Evidence of effectiveness comes from three main sources: (1) the natural experiment of the counteradvertising campaign that occurred during the late 1960s as the result of a Fairness Doctrine ruling (also discussed in "Broadcast Advertising Ban" in Chapter 5), (2) school and community intervention studies incorporating mass media approaches (see "Supplemental Programs" in Chapter 3), and (3) recent experience with large paid

media campaigns in several U.S. states and with a nationwide campaign funded by the FDA. Because of the special sensitivity of young people to tobacco marketing and the high rates of tobacco use among teenagers, the subsequent review in this chapter will focus on countermarketing media campaigns that include prominent youth-targeted components. The literature provides strong evidence of the value of mass media campaigns to inform the public at large—including young people—about the hazards of smoking, to promote specific cessation actions and services (such as telephone help lines), and to provide cessation clinics to adult smokers (Flay 1987; Pierce 1995).

The Fairness Doctrine campaign. In 1967, the Federal Communications Commission (FCC) applied the Fairness Doctrine (discussed in "Broadcast Advertising Ban" in Chapter 5) to cigarette advertising and required broadcasters to provide a significant amount of airtime to antismoking messages—a requirement interpreted by the FCC at that time to be about one antismoking message per three tobacco advertising messages). This requirement resulted in the only sustained nationwide tobacco control media campaign to date. From mid-1967 through 1970, roughly \$200 million in commercial airtime (in 1970 dollars) or \$75 million per year was donated for antismoking messages on television and radio (Warner 1986; USDHHS 1989b).

The campaign produced significant reductions in both adult and youth smoking behaviors (Hamilton 1972). For the first time in the 20th century, adult per capita cigarette consumption fell for more than three consecutive years. Teenage smoking prevalence was 3 percentage points smaller during the Fairness Doctrine period than it was in the 16 months before the campaign, and the campaign was associated overall with a 3.4-percentage point reduction in teen smoking prevalence. Perhaps the ultimate indicator of the campaign's impact was a change that followed the campaign's end: with the 1971 enactment of congressional legislation banning tobacco commercials from television—and with them, the Fairness Doctrine-mandated counteradvertisements—per capita cigarette consumption immediately resumed its upward trend (see "Broadcast Advertising Ban" in Chapter 5).

Hamilton (1972) suggested that during the Fairness Doctrine period, the antismoking campaign messages had an effect that was nearly six times that of cigarette advertisements. Warner (1979) noted that the government's broadcast ban—and the consequent end of the countermarketing campaign—was especially detrimental to the ongoing effort to prevent young

people from smoking. Cigarette promotion remained highly visible in the print media and in tobacco companies' sponsorship of sporting events at the same time the broadcast ban "virtually eliminated mass promotion of the antismoking cause" (p. 445).

Community intervention studies. As described in "Research on Multifaceted Programs" in Chapter 3, multicomponent youth-directed programs that include a prominent mass media component have shown long-term success in postponing or preventing smoking onset in adolescents. In the University of Vermont School and Mass Media Project, the study featuring the most intensive paid counteradvertising campaign, the preventive effect actually increased during the two-year intervention period among the adolescents at higher risk for smoking (Flynn et al. 1997)—a rare outcome for most campaigns trying to change health behaviors. The authors noted that counteradvertising can effectively reach higher-risk youth because of their greater exposure to the mass media, particularly radio and television. It is also likely that higher-risk youth make their decisions about tobacco use earlier in life than lower-risk youth; mass media influences can be especially powerful in shaping attitudes and normative perceptions at early ages.

State-based media campaigns. Mass media campaigns are standard components of the well-funded, ongoing tobacco control programs in California, Massachusetts, Arizona, Florida, and other states receiving money for counteradvertising programs from state excise tax increases or tobacco settlement allotments (as was discussed in "Example of Major State Programs," earlier in this chapter). Although it is difficult to sort out the effectiveness of media campaigns from other program components, evaluations of these statewide public education programs, particularly in California and Massachusetts (see "Supplemental Programs" in Chapter 3), have shown their success in reducing tobacco use among adults, slowing the uptake of tobacco among youth, and protecting children from exposure to ETS (CDC 1996). A recent study of the Massachusetts media campaign in 1993 and 1997 found that among younger adolescents (those aged 12–13 years in 1993), those who had been exposed to the counteradvertising campaign on television were about half as likely to have become smokers as those who had not been able to recall campaign advertisements (Siegel and Biener 2000).

Food and Drug Administration campaign. In 1998, the FDA launched a national advertising campaign to help retailers comply with the age and photo identification provisions of the FDA's rules to prevent tobacco sales to children and adolescents. The

campaign began with a test in Arkansas and by year's end was active in 42 states. Funded annually at about \$9 million, the campaign featured radio spots, billboards, newspaper advertisements, posters, and store signage. The overall approach was to use humor to relieve the discomfort clerks may feel when checking young people's identification/proof-of-age cards and to increase awareness of the rule provisions among retailers, underage youth, and the general population. One counter card, for example, reads, "Our cashier really stinks at guessing ages. So if you want cigarettes, can we see some I.D.?"

A campaign tracking survey (Market Facts 1998) in nine states with test and control sites found that during the first year of the campaign, knowledge of age 27 as the cutoff age for checking identification increased from 34 to 54 percent in test sites and from 31 to 40 percent in control sites. Most important was a small but significant decline in the average number of times minors tried to buy tobacco. According to retailer self-reports, this number declined from 3.4 times each day before the campaign to 2.8 times daily after the media effort. In control sites, the frequency of underage purchase attempts did not decrease from before (2.4 times daily) to after (2.7 times daily) the time of the campaign. For customers from whom identification was requested in the test sites, retailers reported that the proportion of those who were "often" or "always" irritated declined from 34 percent to 28 percent.

Counteradvertising and entertainment media. The increase in movie depictions of tobacco use is a powerful media influence promoting use among teens (Stockwell and Glantz 1997). In focus groups, young people are not able to recall antismoking messages on television or in the movies, but they recall specific movies that portray smoking and can identify actors and actresses who smoke in their entertainment roles (Crawford et al. 1998). Counteradvertising holds promise for helping denormalize and deglamorize these portrayals in the entertainment media. In an experimental study, Pechmann and Shih (1999) found that placement of a 30-second California Department of Health Services tobacco counteradvertisement before the popular movie *Reality Bites* served to inoculate teenagers against the movie's pervasive prosmoking cues without detracting from their enjoyment of the film. Because paid advertising in movie theaters is a highly efficient method of reaching adolescents, the authors recommend this tactic as a nationwide cost-effective prevention strategy.

Research on best practices. Although producers of counteradvertising campaigns use formative research techniques to develop products, inconsistent

testing methods hinder comparison of the effectiveness of different messages. This situation has helped create the impression that there is little agreement over "what works" in tobacco counteradvertising, as typified by this *Washington Post* headline: "The Anti-Smoking Campaign's a Many Splendored Thing, and That's the Problem" (Teinowitz 1998).

Goldman and Glantz (1998), using available focus group data and research reports obtained from a number of states, concluded that two message strategies, industry manipulation and the hazards of ETS, are the most effective for denormalizing smoking among young people and reducing consumption among adults. The researchers reported that addiction and cessation messages can also be effective, but that four strategies are not effective: youth access, short-term health effects, long-term health effects, and romantic rejection. They also characterized California's counteradvertising campaign as more "confrontational with the industry" (p. 772) than Massachusetts' "more youth-oriented approach" (p. 772), citing this difference as a major reason for their finding that the California media campaign was relatively more cost-effective. This paper elicited some strong responses. The University of Vermont School and Mass Media Project investigators (Worden et al. 1998) emphasized the limitations of focus group results and the importance of audience age in reactions to messages. They argued that for young people aged 10 to 12 years (the age group in which they recommended starting prevention efforts), presenting messages that foster positive social influence and social norms have proved most effective in reducing tobacco use among youth. Balch and Rudman (1998) responded that young people participating in 110 focus groups in five different states considered numerous concepts and judged five to be more credible, relevant, and persuasive: addiction, short-term health effects, athletic performance, role model for younger siblings, and effects on family. From Massachusetts, Connolly and Harris (1998) noted that industry manipulation and ETS themes constituted 32 percent of all youth-targeted messages and 37 percent of all messages in the Massachusetts tobacco control media campaign and that on a per capita basis, the state actually outspent California on these messages. Moreover, the researchers reported that Massachusetts experienced a larger decline in per capita cigarette consumption than did California for the period 1990–1996.

To obtain data in a more quantitative way, Pechmann and Shih (1999) created a typology based on 196 youth-oriented antismoking television advertisements. They identified three main types—fear

appeals, peer norms, and tobacco marketing—and further subdivided these into seven main messages: (1) smokers may face serious health problems, (2) tobacco company deception results in disease and death, (3) smokers endanger their family members, (4) smoking is unattractive, (5) smokers are perceived by peers as misguided, (6) most young people choose not to smoke, and (7) advertisement shows how tobacco companies market their products. The investigators tested a sample of 56 of their advertisements in a group of ethnically diverse 7th, 9th, and 10th graders. After viewing a selection of test and placebo advertisements, study participants completed an evaluation survey to assess the effect of each category on their intent to smoke and on other pertinent measures, such as attitudes toward smoking and knowledge of tobacco marketing tactics. Results showed that only three of the seven messages were highly effective in reducing teenagers' intent to smoke: those that conveyed that smokers endanger their family members, that smokers are perceived by peers as misguided, and that most young people choose not to smoke.

In the Massachusetts campaign study (Siegel and Biener 2000), the authors tested eight smoking-related knowledge and attitude variables corresponding to campaign themes. Only one variable, perceived youth smoking prevalence, changed significantly with exposure to the media campaign at baseline and was associated with the reported reduction in tobacco uptake. Exposed youths were more than twice as likely than their unexposed peers to have an accurate perception at follow-up that fewer than half of the students at their high school were smokers. Variables that did not change were knowledge and attitudes related to low-tar cigarettes, environmental tobacco smoke, chemicals, wrinkles, tobacco company tactics, dating, and sports. This finding points to the power of the mass media, especially television, to set social norms and supports the effectiveness of counteradvertising messages that denormalize tobacco use.

As part of a three-year study exploring racial/ethnic and gender differences in teen tobacco use, a group of 11 CDC-funded university-based Prevention Research Centers conducted a series of focus groups during 1996–1997 to explore potentially effective counteradvertising strategies and messages. Six of the 11 centers used television spots from CDC's Media Campaign Resource Center for Tobacco Control to elicit reactions and stimulate discussion. For the most part, different centers used different advertisements, and they did not attempt to "test" the advertisements in any standardized way to determine relative effectiveness. Nevertheless, the conclusions that emerged from

Teen Focus Group Response to Counteradvertising Messages

(Findings from 11 Prevention Research Centers)

- Without an overall context provided by ongoing advertising and other program elements, the message that tobacco companies are manipulating young people to smoke (“they’re lying to you”) has relatively low interest and salience among teens and may be miscomprehended.
- Attempts to explain the concept of nicotine addiction and make it personally relevant for young nonsmokers is difficult because most have not experienced the physical cravings of addiction and tend to take messages literally.
- The television spot shown to the most focus groups (about physical performance and featuring the U.S. Women’s National Soccer Team) was easily understood, attention getting, and credible and may be generalizable (with some effort) to nonathletic endeavors.
- Young people did not like advertisements that feature text.
- Young people, particularly whites, were sharply critical of any advertisement they perceived as corny, “cute,” staged, or unhip.
- As advertising professionals have reported in the research literature, humor was found to be a double-edged sword: it can be very effective, but if used inappropriately can be seen as trivializing the issue. In some focus groups, humorous advertisements obtained both the highest and the lowest scores.
- Young people reacted emotionally and favorably to true, nonpreachy stories about the impact of smoking on a person’s or family member’s life (such as a television spot from California featuring a man whose wife had died from exposure to his smoking).
- Cartoons tend to have low “stopping power” because teens have seen so many, whereas the use of surprising characters like animals (such as the “Animals” and “Butts” spots from Minnesota) can rivet attention. These attention-getting spots do not necessarily communicate an effective countermessage, however.
- Messages that portray the negative social effects of tobacco use perform well among teens; messages that focus on health effects can be effective if they are presented dramatically but realistically (such as a California spot featuring a laryngectomy patient smoking a cigarette).

this research (Tobacco Network, unpublished data) give some indication of the complexity of people’s response and the considerable challenges to crafting effective messages (see the text box “Teen Focus Group Response to Counteradvertising Messages”).

Audience targeting. The use of counteradvertising aimed only at young people rather than the use of a general marketing approach has been controversial. Glantz (1996) criticized the public health community’s “preoccupation with youth” (p. 157), particularly youth access campaigns, as an ineffective strategy and one that diverts energy from reducing adult smoking and creating a smoke-free society. Cummings and Clarke (1998) warned that campaigns focused exclusively on young people may be counterproductive if the messages make smoking more appealing to youth by promoting it as something that is

not for them. Indeed, a chief criticism of the tobacco industry-funded booklet *Tobacco: Helping Youth Say No* was that it portrayed tobacco use as a forbidden fruit and a badge of maturity, thereby increasing its attraction to youth (DiFranza and McAfee 1992). The Institute of Medicine noted that “as adolescents venture more and more into the community, their perceptions that certain norms seem to apply only to them and not to adults may promote health-compromising behaviors” (Lynch and Bonnie 1994, p. 87). Young people participating in focus groups conducted during the third year (1997–1998) of the CDC-funded Tobacco Network project reported that they respect and regard policies targeted to the public at large, such as clean indoor air laws, but resent policies specific to them, such as youth access restrictions. They also resented the inconsistent enforcement of general

Tips for Success in Health Promotion Campaigns

- **Target young people in grades six and nine (ages 11 and 15).** These years define critical periods in most children's social development, times when many young people change schools and peer groups.
- **Target adults with complementary, noncontradictory messages.** In a comprehensive strategy, media messages that inevitably spill over from one audience to another can be mutually reinforcing and synergistic. Clean indoor air messages can provide added motivation for adults to quit smoking. Cessation messages for adults can affect young people's perception of norms and highlight the problem of addiction. Prevention messages for young people can increase the salience of the tobacco issue among parents and community leaders.
- **Highlight nonsmoking as the majority behavior.** Most young people overestimate the number of their peers who use tobacco. Campaigns should not seek to correct this misperception and highlight an increasing "problem" of kids who smoke.
- **Present realistic tobacco-free lifestyles** as practiced by diverse, appealing, and interesting persons. Youth behaviors are driven by how young people perceive the behaviors of people like them. Having a repertoire of social choices is a fundamental need for teens, who are going through a period of profound social and environmental transition.
- **Provide constructive alternatives** to tobacco use and discourage destructive alternatives. Sports and other youth-oriented activities associated with the tobacco-free lifestyle can provide some of that positive social repertoire.
- **Communicate the relevant dangers** of tobacco. Certain dangers of tobacco, if explained in a creative and memorable manner, resonate with young people—for example, addiction portrayed as a loss of control, the carcinogenicity of environmental tobacco smoke, the toxic chemicals in tobacco products and smoke, and the tangible suffering and visible disfigurement from tobacco-related diseases. Communicate health messages through personal testimonies (tell a story) and creative executions that break through young people's sense of immortality and their (and adults') resistance to traditional health messages.
- **Encourage youth empowerment and control.** Teens need to be offered information and anecdotal experience from which they can begin to understand the world and take control of their own lives.
- **Abandon the search for the "magic-bullet" message.** There is no single best motivator for preventing or reducing tobacco use. Campaign messages for both young people and adults should feature a variety of themes, appeals (fear, humor, satire, testimonials, etc.), and executional styles. Maximize the number, variety, and novelty of messages rather than communicating a few messages repeatedly.

continued on next page

policies, such as allowing teachers but not students to smoke on school property.

Worden (in Cummings and Clarke 1998), referring to the research literature on multifaceted education campaigns, noted that reducing the demand for tobacco among young people requires a combination of direct (to youth) and indirect (to adults) messages and careful attention to audience segmentation. He stressed that young people and adults need separate media campaigns that do not contradict each other. For example, a youth-directed television spot that communicates the

message "most kids don't smoke" can be neutralized by an adult-aimed but youth-viewed spot that says "more and more kids are smoking every day."

Characteristics of Successful Campaigns

Though debate continues over the relative effectiveness of strategies employing specific messages, the experience reviewed in preceding sections suggests consensus that counteradvertising campaigns must have sufficient reach, frequency, and duration to be

continued

- **Use multiple nonpreachy voices.** Not only do different teens require different appeals and creative executions, but diversity of messages is itself a sophisticated message. Teens strongly reject attempts by anyone to dominate or direct them. Messages about industry manipulation, if they are to be relevant and acceptable to youth, should be delivered by nonauthoritarian sources (such as Florida's "Truth" campaign teenagers), not with melodramatic appeals. Avoid highlighting a single theme, tagline, identifier, or sponsor.
- **Use a complementary, reinforcing mix of television, radio, print, and outdoor advertising.** The campaign should also explore the various alternative media options available (e.g., movie trailers, the Internet, other computer resources, video games, materials for schools and community groups). The media mix is especially important in view of today's proliferating fragmented media market.
- **Involve parents and families** in activities that will reduce risk factors and promote protective factors for young people at risk for tobacco use. Parents and other family members have substantial influence on the perceptions and behaviors of young people.
- **Maximize use of existing high-quality media materials** produced by the government, voluntary agencies, and a number of individual states. (A new, high-quality television spot commonly costs more than \$100,000 to produce.) A large collection of advertisements is currently available through the CDC's Media Campaign Resource Center for Tobacco Control. The cost of placing an advertisement will vary significantly by state and media market.
- **Include grassroots promotions, local media advocacy, event sponsorships, and other community tie-ins** to support and reinforce the counteradvertising campaign (see "Media Advocacy," earlier in this chapter). Work in concert with other interventions to promote policies that aim to change social norms regarding tobacco. A local "look" for local media messages (e.g., featuring people of ethnic or geographic representation similar to the viewing audience) appears to be more important for adults than for youth, because young people tend to share and be shaped by a more universal, multiethnic youth "media world."

successful. The 1967–1970 media campaign, enabled by the Fairness Doctrine, achieved high frequency (one antismoking advertisement per three cigarette advertisements), extended reach (virtually complete audience penetration through three [pre-cable television] national networks), and long duration (three and a half years). The youth-aimed media campaign of the University of Vermont School and Mass Media Project exposed 50 percent of the target population to each television and radio spot about 6 times each year over a four-year period (about the midpoint in the recommended exposure range of 3 to 10 times per year). This level of exposure is possible only through paid media placement.

Another lesson from health promotion campaigns is the need for research at every phase of campaign planning and implementation. Campaigns should be grounded in the extensive literature on psychosocial risk factors for initiating, continuing, and stopping tobacco use and should be guided by expertise in

communications theory and practice. Media materials should undergo rigorous audience pretesting to ensure they achieve predetermined communication objectives with their target audiences. Ongoing measurement of the communications' impact is needed to evaluate the effectiveness of the campaign and to guide midcourse corrections.

Through the Columbia University Prevention Research Center in New York City, the CDC convened a panel of youth marketing and research experts in 1996 to advise the agency on effective countermarketing approaches to prevent tobacco use among young people. Over two years, the expert panel reviewed the literature, interviewed experts in tobacco control and health promotion, and drew on their private-sector experience and resources to develop a set of strategic guidelines for such a campaign (McKenna et al. 2000). This work, supplemented by other reviews of counteradvertising campaigns (USDHHS 1994; Pechmann 1997; Siegel 1998; Teenage Research

Unlimited 1999; Pechmann and Reibling 2000), yielded recommendations for effective media campaigns to prevent tobacco use (see the text box "Tips for Success in Health Promotion Campaigns").

These recommendations serve as general guidance for tobacco counteradvertising efforts, but further research is needed to refine our understanding of the role and effects of mass media. Relevant areas for further investigation include determining the impact of

counteradvertising on tobacco use behaviors, on readiness to quit, on attitudes toward tobacco advertising and tobacco use, and on other predictors of initiation and cessation; identifying the most effective themes, techniques, and messages; tailoring messages to high-risk groups; exploring the role of new communication tools, such as the Internet; attributing impact; and examining the interaction of media campaigns with private and public tobacco control policies.

Summary

The conceptual framework described at the start of this chapter defines the basic components of the health promotion intervention model. The statewide tobacco control programs being funded either by increases in cigarette excise taxes or settlements with the tobacco industry are creating a new laboratory to test many of these conceptual models for comprehensive tobacco control. Recently, both the Institute of Medicine (IOM) and researchers have released reviews of the emerging data from these statewide tobacco control efforts. In their report, the IOM (2000) noted that "it is difficult to attribute a reduction in tobacco use to any single factor; nevertheless, they conclude that "multifaceted state tobacco control programs are effective in reducing tobacco use" (p. 4). In a review focusing more specifically on the effectiveness of these new statewide tobacco control programs on teenage smoking, Wakefield and Chaloupka (1999) conclude that "There is consistent evidence the programs are associated with a decline in adult smoking prevalence" (p. 6), but they are somewhat more cautious about the impact of these programs on youth smoking. Nevertheless, they do conclude that "Notwithstanding these cautions, we find that the weight of evidence falls in favor of comprehensive tobacco control programs being able to reduce teenage tobacco use" (p. 6).

In the consideration of the emerging data from these statewide tobacco control programs, it is important to note that many programmatic elements of the comprehensive tobacco control program framework are still being refined and evaluated. Thus, no current statewide program serves an ideal or model program. Wakefield and Chaloupka (1999) conducted a careful review of the various elements of the statewide

programs in Arizona, California, Florida, Massachusetts, and Oregon. They placed special attention on the strengths of the "inputs"—"namely, what was *actually implemented* as part of the programs." Additionally, they assessed how "actual implementation of program strategies may differ substantially from intended implementation" and noted that "the extent of disparity may vary over time and between programs." Much more evaluation research is needed in order to sort out the efficacy of individual components of these evolving comprehensive programs and to refine the comprehensive program structure.

Finally, although the data from these statewide tobacco control programs are encouraging, these results need to be considered in the perspective of the less favorable results from the community trials. The conceptual framework for the comprehensive tobacco control programs shares many elements with the theoretical models used to develop the community trial interventions. However, as Wakefield and Chaloupka (1999) noted, the programs actually implemented may differ substantially from the intended implementation. There has been some effort to analyze how the program components within the emerging statewide tobacco control programs may differ from interventions tested within the community trials (Green and Richard 1993; Schmid et al. 1995), but much more work is needed in this area. As the IOM (2000) and Wakefield and Chaloupka (1999) concluded, the results from the statewide tobacco control programs are favorable. However, both reviews emphasize the importance of continued surveillance and evaluation efforts to monitor program performance, to provide accountability for the use of public funds, and to improve program efforts.

Conclusions

1. The large-scale interventions conducted in community trials have not demonstrated a conclusive impact on preventing and reducing tobacco use.
2. Statewide programs have emerged as the new laboratory for developing and evaluating comprehensive plans to reduce tobacco use.
3. Initial results from the statewide tobacco control programs are favorable, especially regarding declines in per capita consumption of tobacco products.
4. Results of statewide tobacco control programs suggest that youth behaviors regarding tobacco use are more difficult to change than adult ones, but initial results of these programs are generally favorable.

References

- Abramson JH, Gofin R, Hopp C, Gofin J, Donchin M, Habib J. Evaluation of a community program for the control of cardiovascular risk factors: the CHAD program in Jerusalem. *Israel Journal of Medical Sciences* 1981;17(2-3):201-12.
- Abt Associates Inc. for the Massachusetts Department of Public Health. *Independent Evaluation of the Massachusetts Tobacco Control Program*. First Annual Report: Fiscal Year 1994. Cambridge (MA): Abt Associates Inc., 1995.
- Abt Associates Inc. for the Massachusetts Department of Public Health. *Independent Evaluation of the Massachusetts Tobacco Control Program*. Fourth Annual Report. Cambridge (MA): Abt Associates Inc., 1997.
- Abt Associates Inc. for the Massachusetts Department of Public Health. *Independent Evaluation of the Massachusetts Tobacco Control Program*. Fifth Annual Report: Summary. Cambridge (MA): Abt Associates Inc., 1999.
- Advocacy Institute. *Philip Morris Corporate Image Advertising Campaign*. Action Alert. Washington: Advocacy Institute, Smoking Control Advocacy Resource Center (SCARC), Nov 3, 1989.
- Advocacy Institute. *The Smoking Control Advocacy Resource Center Network: Evaluation of SCARCNet*. Washington: Advocacy Institute, 1994.
- Allison KR, Rootman I. Scientific rigor and community participation in health promotion research: are they compatible? *Health Promotion International* 1996; 11(4):333-40.
- Altman DG, Flora JA, Fortmann SP, Farquhar JW. The cost-effectiveness of three smoking cessation programs. *American Journal of Public Health* 1987;77(2): 162-5.
- American Medical Association. *SmokeLess States National Tobacco Prevention and Control Program: Five Years of Making a Difference*. Chicago: American Medical Association, 1998.
- American Medical News*. No more Dakotas. *American Medical News*, Apr 13, 1992; Roundup Sect:18 (col 3).
- American Public Health Association. *International Network of Women Against Tobacco Member Directory* 1994. Washington: American Public Health Association, 1994.
- Americans for Nonsmokers' Rights v. The State of California*, 51 Cal. App. 4th 743, 1996 Cal. App. LEXIS 1165, 59 Cal. (Cal. Ct. App., Third App. Dist. 1996).
- Anantha N, Nandakumar A, Vishwanath N, Venkatesh T, Pallad YG, Manjunath P, Kumar DR, Murthy SGS, Shivashankariah, Dayananda CS. Efficacy of an anti-tobacco community education program in India. *Cancer Causes and Control* 1995;6(2):119-29.
- Aravanis C, Corcondilas A, Dontas AS, Lekos D, Keys A. Coronary heart disease in seven countries. IX. The Greek islands of Crete and Corfu. *Circulation* 1970;41(4 Suppl):I88-I100.
- Arizona Cancer Center, Behavioral Sciences Section, for the Arizona Department of Health Services. *Evaluation of the Tobacco Education and Prevention Program Media Campaign*. Tucson (AZ): Arizona Cancer Center, 1998.
- Arizona Department of Health Services. *1999 Arizona Adult Tobacco Survey Report*. Phoenix (AZ): Arizona Department of Health Services, Bureau of Public Health Statistics, 2000.
- Bal DG, Kizer KW, Felten MA, Mozar HN, Niemeyer D. Reducing tobacco consumption in California: development of a statewide anti-tobacco use campaign. *Journal of the American Medical Association* 1990;264(12): 1570-4.
- Balch GI, Rudman G. Antismoking advertising campaigns for youth [letter]. *Journal of the American Medical Association* 1998;280(4):323-4.
- Bandura A. *Social Learning Theory*. Englewood Cliffs (NJ): Prentice-Hall, 1977.
- Bauer RL, Heller RF, Challah S. United Kingdom Heart Disease Prevention Project: 12-year follow-up of risk factors. *American Journal of Epidemiology* 1985;121(4): 563-9.

Baum F. Researching public health: behind the qualitative–quantitative methodological debate. *Social Science and Medicine* 1995;40(4):459–68.

Begay ME, Traynor M, Glantz SA. The tobacco industry, state politics, and tobacco education in California. *American Journal of Public Health* 1993;83(9):1214–21.

Biener L, Siegel M. Tobacco marketing and adolescent smoking: more support for a causal inference. *American Journal of Public Health* 2000;90(3):407–11.

Blackburn H. Research and demonstration projects in community cardiovascular disease prevention. *Journal of Public Health Policy* 1983;4(4):398–421.

Blackburn H, Taylor HL, Keys A. Coronary heart disease in seven countries. XVI. The electrocardiogram in prediction of five-year coronary heart disease incidence among men aged forty through fifty-nine. *Circulation* 1970;41(4 Suppl):I154–I61.

Blum A. Medical activism. In: Taylor RB, Ureda JR, Denham JW, editors. *Health Promotion: Principles and Clinical Applications*. Norwalk (CT): Appleton-Century-Crofts, 1982:373–91.

Bracht N, Finnegan JR Jr, Rissel C, Weisbrod R, Gleason J, Corbett J, Veblen-Mortenson S. Community ownership and program continuation following a health demonstration project. *Health Education Research* 1994; 9(2):243–55.

Briton NJ, Clark TW, Baker AK, Posner J, Soldz S, Krakow M. *Adolescent Tobacco Use in Massachusetts. Trends Among Public School Students 1984–1996*. Boston: Health and Addictions Research, 1997.

Butler J. The “Dakota Papers.” Paper presented at the Seventh World Conference on Tobacco and Health; Mar 30, 1990; Perth (Australia).

Buzina R, Keys A, Mohacek I, Marinkovic M, Hahn A, Blackburn H. Coronary heart disease in seven countries. V. Five-year follow-up in Dalmatia and Slavonia. *Circulation* 1970;41(4 Suppl):I40–I51.

Carleton RA, Lasater TM, Assaf AR, Feldman HA, McKinlay S, and the Pawtucket Heart Health Program Writing Group. The Pawtucket Heart Health Program: community changes in cardiovascular risk factors and projected disease risk. *American Journal of Public Health* 1995;85(6):777–85.

Centers for Disease Control. State coalitions for prevention and control of tobacco use. *Morbidity and Mortality Weekly Report* 1990;39(28):476–84.

Centers for Disease Control and Prevention. Changes in the cigarette brand preferences of adolescent smokers—United States, 1989–1993. *Morbidity and Mortality Weekly Report* 1994; 43(32):577–81.

Centers for Disease Control and Prevention. Cigarette smoking before and after an excise tax increase and an antismoking campaign—Massachusetts, 1990–1996. *Morbidity and Mortality Weekly Report* 1996;45(44): 966–70.

Centers for Disease Control and Prevention. Tobacco tax initiative—Oregon, 1996. *Morbidity and Mortality Weekly Report* 1997;46(11):246–8.

Centers for Disease Control and Prevention. *1998 Program Briefing Book*. Atlanta: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1998.

Centers for Disease Control and Prevention. *Best Practices for Comprehensive Tobacco Control Programs—August 1999*. Atlanta: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1999a.

Centers for Disease Control and Prevention. Decline in cigarette consumption following implementation of a comprehensive tobacco prevention and education program—Oregon, 1996–1998. *Morbidity and Mortality Weekly Report* 1999b;48(7):140–3.

Centers for Disease Control and Prevention. Tobacco use among middle and high school students—Florida, 1998 and 1999. *Morbidity and Mortality Weekly Report* 1999c;48(12):248–53.

Chaloupka FJ, Grossman M. *Price, tobacco control policies and youth smoking*. Working paper no. 5740. Cambridge (MA): National Bureau of Economic Research, 1996.

COMMIT Research Group. Community Intervention Trial for Smoking Cessation (COMMIT): summary of design and intervention. *Journal of the National Cancer Institute* 1991;83(22):1620–8.

COMMIT Research Group. Community Intervention Trial for Smoking Cessation (COMMIT): I. Cohort results from a four-year community intervention. *American Journal of Public Health* 1995a;85(2):183-92.

COMMIT Research Group. Community Intervention Trial for Smoking Cessation (COMMIT): II. Changes in adult cigarette smoking prevalence. *American Journal of Public Health* 1995b;85(2):193-200.

Connolly GN, Harris JE. Evaluating antismoking advertising campaigns [letter]. *Journal of the American Medical Association* 1998;280(11):964-5.

Crawford M, Headen S, Luke D, McGloin T. Latest research: youth and teens. Presented at the National Conference on Tobacco and Health; Oct 26-28, 1998; St. Paul (MN).

Cummings KM, Clarke H. *The Use of Counter-Advertising as a Tobacco Use Deterrent and Analysis of Pending Federal Tobacco Legislation*. Policy Analysis No. 8. Washington: Advocacy Institute, Health Science Analysis Project, 1998.

DiFranza JR, McAfee T. The Tobacco Institute: helping youth say "yes" to tobacco. *Journal of Family Practice* 1992;34(6):694-6.

DiFranza JR, Richards JW, Paulman PM, Wolf-Gillespie N, Fletcher C, Jaffe RD, Murray D. RJR Nabisco's cartoon camel promotes Camel cigarettes to children. *Journal of the American Medical Association* 1991;266(22):3149-53.

Downie RS, Tannahill C, Tannahill A. *Health Promotion: Models and Values*. 2nd ed. Oxford: Oxford University Press, 1996.

Dwyer T, Pierce JP, Hannam CD, Burke N. Evaluation of the Sydney "Quit. For Life" anti-smoking campaign. Part 2: changes in smoking prevalence. *Medical Journal of Australia* 1986;144(7):344-7.

Egger G, Fitzgerald W, Frape G, Monaem A, Rubinstein P, Tyler C, McKay B. Results of large scale media anti-smoking campaign in Australia: North Coast "Quit for Life" programme. *British Medical Journal* 1983;287(6399):1125-8.

Elder JP, McGraw SA, Rodrigues A, Lasater TM, Ferreira A, Kendall L, Peterson G, Carleton RA. Evaluation of two community-wide smoking cessation contests. *Preventive Medicine* 1987;16(2):221-34.

Elder JP, Schmid TL, Dower P, Hedlund S. Community heart health programs: components, rationale, and strategies for effective interventions. *Journal of Public Health Policy* 1993;14(4):463-79.

Eriksen MP. Social forces and tobacco in society. Paper presented at the Robert Wood Johnson Foundation Conference, New Partnerships and Paradigms for Tobacco Prevention Research; May 6-9, 1997; Sundance (UT).

Farquhar JW. The community-based model of life style intervention trials. *American Journal of Epidemiology* 1978;108(2):103-11.

Farquhar JW, Fortmann SP, Flora JA, Taylor B, Haskell WL, Williams PT, Maccoby N, Wood PD. Effects of communitywide education on cardiovascular disease risk factors: the Stanford Five-City Project. *Journal of the American Medical Association* 1990;264(3):359-65.

Farquhar JW, Fortmann SP, Maccoby N, Haskell WL, Williams PT, Flora JA, Taylor CB, Brown BW Jr, Solomon DS, Hulley SB. The Stanford Five-City Project: design and methods. *American Journal of Epidemiology* 1985;122(2):323-34.

Farquhar JW, Maccoby N, Wood PD, Alexander JK, Breitrose H, Brown BW Jr, Haskell WL, McAlister AL, Meyer AJ, Nash JD, Stern MP. Community education for cardiovascular health. *Lancet* 1977;1(8023):1192-5.

Farquhar JW, Magnus PF, Maccoby N. The role of public information and education in cigarette smoking controls. *Canadian Journal of Public Health* 1981;72(6):412-20.

Federal Register. US Department of Health and Human Services, Food and Drug Administration. Regulations restricting the sale and distribution of cigarettes and smokeless tobacco to protect children and adolescents (21 CFR Parts 801, 803, 804, 807, 820, and 897), 61 *Fed. Reg.* 44396 (1996).

Fidanza F, Puddu V, Imbimbo AB, Menotti A, Keys A. Coronary heart disease in seven countries. VII. Five-year experience in rural Italy. *Circulation* 1970;41(4 Suppl): I63-I75.

Fisher EB Jr. The results of the COMMIT trial [editorial]. *American Journal of Public Health* 1995;85(2):159–60.

Fisher EB, Auslander WF, Munro JF, Arfken CL, Brownson RC, Owens NW. Neighbors for a Smoke Free North Side: evaluation of a community organization approach to promoting smoking cessation among African Americans. *American Journal of Public Health* 1998;88(11):1658–63.

Flay BR. *Selling the Smokeless Society: Fifty-six Evaluated Mass Media Programs and Campaigns Worldwide*. Washington: American Public Health Association, 1987.

Florida Department of Health. 2000 Florida Youth Tobacco Survey results. *Florida Youth Tobacco Survey* 2000;3(1).

Flynn BS, Worden JK, Secker-Walker RH, Pirie PL, Badger GJ, Carpenter JH. Long-term responses of higher and lower risk youths to smoking prevention interventions. *Preventive Medicine* 1997;26(3):389–94.

Fortmann SP, Taylor CB, Flora JA, Jatulis DE. Changes in adult cigarette smoking prevalence after 5 years of community health education: the Stanford Five-City Project. *American Journal of Epidemiology* 1993;137(1):82–96.

Gail MH, Byar DP, Pechacek TF, Corle DK, for the COMMIT Study Group. Aspects of statistical design for the Community Intervention Trial for Smoking Cessation (COMMIT). *Controlled Clinical Trials* 1992;13(1):6–21.

German Cardiovascular Prevention Study Group. The German Cardiovascular Prevention Study (GCP): design and methods. *European Heart Journal* 1988;9(10):1058–66.

Giovino GA, Schooley MW, Zhu B-P, Chrismon JH, Tomar SL, Peddicord JP, Merritt RK, Husten CG, Eriksen MP. Surveillance for selected tobacco-use behaviors—United States, 1900–1994. *Morbidity and Mortality Weekly Report* 1994;43(SS-3):1–43.

Glantz SA. Achieving a smokefree society [editorial]. *Circulation* 1987;76(4):746–52.

Glantz, SA. Preventing tobacco use—the youth access trap [editorial]. *American Journal of Public Health* 1996;86(2):156–8.

Glantz SA, Begay ME. Tobacco industry campaign contributions are affecting tobacco control policymaking in California. *Journal of the American Medical Association* 1994;272(15):1176–82.

Glasgow RE, Cummings KM, Hyland A. Relationship of worksite smoking policy to changes in employee tobacco use: findings from COMMIT. Community Intervention Trial for Smoking Cessation. *Tobacco Control* 1997;6(2 Suppl):S44–8.

Glasgow RE, Klesges RC, Mizes JS, Pechacek TF. Quitting smoking: strategies used and variables associated with success in a stop-smoking contest. *Journal of Consulting and Clinical Psychology* 1985;53(6):905–12.

Gofin J, Gofin R, Abramson JH, Ban R. Ten-year evaluation of hypertension, overweight, cholesterol, and smoking control: the CHAD program in Jerusalem. *Preventive Medicine* 1986;15(3):304–12.

Goldman LK, Glantz SA. Evaluation of antismoking advertising campaigns. *Journal of the American Medical Association* 1998;279(10):772–7.

Goodenow C. Tobacco Use Among High School Students: 1999 Massachusetts Youth Risk Behavior Survey Results, Massachusetts Department of Education; <<http://www.doe.mass.edu/news/november99/112299yrb.html>>; accessed June 6, 2000.

Green LW, Kreuter MW. *Health Promotion Planning: An Educational and Environmental Approach*. 2nd ed. Mountain View (CA): Mayfield Publishing Company, 1991.

Green LW, Richard L. The need to combine health education and health promotion: the case of cardiovascular disease prevention. *Promotion and Education* Dec 1993;11–7.

Group Against Smokers' Pollution. *Non-Smokers' Bill of Rights*. College Park (MD): Group Against Smokers' Pollution, n.d.

Gutzwiller F, Nater B, Martin J. Community-based primary prevention of cardiovascular disease in Switzerland: methods and results of the National Research Program (NRP 1A). *Preventive Medicine* 1985;14(4):482–91.

Hamilton JL. The demand for cigarettes: advertising, the health scare, and the cigarette advertising ban. *Review of Economics and Statistics* 1972;54(4):401–11.

Health Promotion International. The Jakarta Declaration on leading health promotion into the 21st century. *Health Promotion International* 1997;12(4):261-4.

Heller K. The slow burn: cigarettes, cancer and blacks. *The Philadelphia Inquirer Magazine*, May 20, 1990;14, 18-20, 30, 32-34, 36.

Hoffmeister H, Mensink GBM, Stolzenberg H, Hoeltz J, Kreuter H, Laaser U, Nüssel E, Hüllemann K-D, v. Troschke J. Reduction of coronary heart disease risk factors in the German Cardiovascular Prevention Study. *Preventive Medicine* 1996;25(2):135-45.

Hu T-w, Sung HY, Keeler TE. Reducing cigarette consumption in California: tobacco taxes vs an anti-smoking media campaign. *American Journal of Public Health* 1995;85(9):1218-22.

Independent Evaluation Consortium. *Final Report of the Independent Evaluation of the California Tobacco Control Prevention and Education Program: Wave I Data, 1996-1997*. Rockville (MD): The Gallup Organization, 1998.

Institute of Medicine. *State Programs Can Reduce Tobacco Use*. A Report of the National Cancer Policy Board. Washington: National Research Council, 2000.

International Network of Women Against Tobacco. Herstories. *World Smoking & Health* 1994;19(2):1-16.

Interreligious Coalition on Smoking OR Health. *Purpose*. Takoma Park (MD): Interreligious Coalition on Smoking OR Health, 1993.

Interreligious Coalition on Smoking OR Health. Honor roll. *Religion & Tobacco Control* Jan 1994:1.

Jacobs DR Jr, Luepker RV, Mittelmark MB, Folsom AR, Pirie PL, Mascioli SR, Hannan PJ, Pechacek TF, Bracht NE, Carlaw RW, et al. Community-wide prevention strategies: evaluation of the Minnesota Heart Health Program. *Journal of Chronic Diseases* 1986;39(10):775-88.

Jacobson B. Smoking and health: a new generation of campaigners. *British Medical Journal* 1983;287(6390):483-4.

Kann L, Kinchen SA, Williams BI, Ross JG, Lowry R, Hill CV, Grunbaum JA, Blumson PS, Collins JL, Kolbe LJ. Youth risk behavior surveillance—United States, 1997. *Morbidity and Mortality Weekly Report* 1998;47(SS-3):1-89.

Kickbusch I. Approaches to an ecological base for public health. *Health Promotion* 1989;4(4):265-8.

Kimura N, Keys A. Coronary heart disease in seven countries. X. Rural southern Japan. *Circulation* 1970;41(4 Suppl):I101-I112.

King AC, Flora JA, Fortmann SP, Taylor CB. Smokers' challenge: immediate and long-term findings of a community smoking cessation contest. *American Journal of Public Health* 1987;77(10):1340-1.

Korhonen HJ, Niemensivu H, Piha T, Koskela K, Wiio J, Johnson CA, Puska P. National TV smoking cessation program and contest in Finland. *Preventive Medicine* 1992;21(1):74-87.

Korhonen HJ, Puska P, Lipand A, Kasmel A. Combining mass media and contest in smoking cessation: an experience from a series of national activities in Finland. *Hygie* 1993;12(1):14-8.

Kornitzer M, Dramaix M, Kittel F, De Backer G. The Belgian Heart Disease Prevention Project: changes in smoking habits after two years of intervention. *Preventive Medicine* 1980;9(4):496-503.

Koskela K. *A Community-Based Antismoking Programme as a Part of a Comprehensive Cardiovascular Programme (The North Karelia Project)*. Kuopio (Finland): Research Institute of Public Health, University of Kuopio, 1981.

Lando HA, Loken B, Howard-Pitney B, Pechacek T. Community impact of a localized smoking cessation contest. *American Journal of Public Health* 1990;80(5):601-3.

Lando HA, Pechacek TF, Pirie PL, Murray DM, Mittelmark MB, Lichtenstein E, Nothwehr F, Gray C. Changes in adult cigarette smoking in the Minnesota Heart Health Program. *American Journal of Public Health* 1995;85(2):201-8.

Lando HA, Pirie PL, McGovern PG, Pechacek TF, Swim J, Loken B. A comparison of self-help approaches to smoking cessation. *Addictive Behaviors* 1991;16(5):183-93.

- Lefebvre RC, Lasater TM, Carleton FA, Peterson G. Theory and delivery of health programming in the community: the Pawtucket Heart Health Program. *Preventive Medicine* 1987;16(1):80-95.
- Lichtenstein E, Thompson B, Nettekoven L, Corbett K for the COMMIT Research Group. Durability of tobacco control activities in 11 North American communities: life after the Community Intervention Trial for Smoking Cessation (COMMIT). *Health Education Research* 1996;11(4):527-34.
- Lichtenstein E, Wallack L, Pechacek TF for the COMMIT Research Group. Introduction to the Community Intervention Trial for Smoking Cessation (COMMIT). *International Quarterly of Community Health Education* 1990-91;11(3):173-85.
- Luepker RV. Community trials. *Preventive Medicine* 1994;23(5):602-5.
- Luepker RV, Murray DM, Jacobs DR Jr, Mittelmark MB, Bracht N, Carlaw R, Crow R, Elmer P, Finnegan J, Folsom AR, Grimm R, Hannan PJ, Jeffrey R, Lando H, McGovern P, Mullis R, Perry CL, Pechacek T, Pirie P, Sprafka JM, Weisbrod R, Blackburn H. Community education for cardiovascular disease prevention: risk factor changes in the Minnesota Heart Health Program. *American Journal of Public Health* 1994;84(9):1383-93.
- Lynch BS, Bonnie RJ, editors. *Growing Up Tobacco Free: Preventing Nicotine Addiction in Children and Youths*. Washington: National Academy Press, 1994.
- Macaskill P, Pierce JP, Simpson JM, Lyle DM. Mass media-led antismoking campaign can remove the education gap in quitting behavior. *American Journal of Public Health* 1992;82(1):96-8.
- Maccoby N, Farquhar JW, Wood PD, Alexander J. Reducing the risk of cardiovascular disease: effects of a community-based campaign on knowledge and behavior. *Journal of Community Health* 1977;3(2):100-14.
- Macdonald G, Veen C, Tones K. Evidence for success in health promotion: suggestions for improvement. *Health Education Research* 1996;11(3):367-76.
- Madonna MG. Getting key players to work together and defending against diversion—Arizona. *Cancer* 1998;83(12 Suppl):2708-12.
- Manley M, Lynn W, Epps RP, Grande D, Glynn T, Shopland D. The American Stop Smoking Intervention Study for cancer prevention: an overview. *Tobacco Control* 1997a;6(Suppl 2):S5-S11.
- Manley MW, Pierce JP, Gilpin EA, Rosbrook B, Berry C, Wun L-M. Impact of the American Stop Smoking Intervention Study on cigarette consumption. *Tobacco Control* 1997b;6(Suppl 2):S12-S16.
- Market Facts. *Tobacco Retailer Study*. Prepared for the Food and Drug Administration, 1998. Boston: Arnold Communications, 1998.
- McAlister A, Puska P, Salonen JT, Tuomilehto J, Koskela K. Theory and action for health promotion: illustrations from the North Karelia Project. *American Journal of Public Health* 1982;72(1):43-50.
- McAlister AL, Farquhar JW, Thoresen CE, Maccoby N. Behavioral science applied to cardiovascular health: progress and research needs in the modification of risk-taking habits in adult populations. *Health Education Monographs* 1976;4(1):45-74.
- McGuire WJ. Persuasion, resistance, and attitude change. In: Pool I, Frey FW, Schramm W, Maccoby N, Parker EB, editors. *Handbook of Communication*. Chicago: Rand McNally College Publishing Company;1973: 216-52.
- McKenna J, Gutierrez K, McCall K. Strategies for an effective youth counter-marketing program: recommendations from commercial marketing experts. *Journal of Public Health Management Practice* 2000;6(3):7-13.
- McMillen RC, Ritchie LA, Robertson AA, Frese W, Cosby AG. *The 1999 Mississippi Social Climate Survey of Tobacco Control and Tobacco Use: Establishing a Benchmark for Mississippi. A Report to the Mississippi State Department of Health*. Mississippi State (MS): Mississippi State University, Social Science and Research Center, 1999.
- Meister JS. Designing an effective statewide tobacco control program—Arizona. *Cancer* 1998;83(12 Suppl): 2728-32.
- Meyer AJ, Nash JD, McAlister AL, Maccoby N, Farquhar JW. Skills training in a cardiovascular health education campaign. *Journal of Consulting and Clinical Psychology* 1980;48(2):129-42.

Minnesota Department of Health. *The Minnesota Plan for Nonsmoking and Health: Report and Recommendations of the Technical Advisory Committee on Nonsmoking and Health*. Minneapolis: Minnesota Department of Health, 1984.

Minnesota Department of Health. *The Minnesota Tobacco-Use Prevention Initiative January 1987–December 1988: A Report to the 1989 Legislature*. Minneapolis: Minnesota Department of Health, 1989.

Minnesota Department of Health. *Minnesota Tobacco-Use Prevention Initiative 1989–1990: A Report to the 1991 Legislature*. Minneapolis: Minnesota Department of Health, 1991.

Minnesota Department of Health. *Minnesota Youth Tobacco Prevention Initiative: Strategic Plan, September 1999–July 2001*. St. Paul (MN): Minnesota Department of Health, 1999.

Mintz M. Where there's smoke. *The Washington Post Magazine*, Dec 3, 1995:20–8, 46–8.

Mississippi State Department of Health. *Mississippi's 1999 Youth Risk Behavior Survey: Summary of Findings*. Jackson (MS): Mississippi State Department of Health, 2000.

Mittelmark MB, Hunt MK, Heath GW, Schmid TL. Realistic outcomes: lessons from community-based research and demonstration programs for the prevention of cardiovascular diseases. *Journal of Public Health Policy* 1993;14(4):437–62.

Moon RW, Males MA, Nelson DE. The 1990 Montana initiative to increase cigarette taxes: lessons for other states and localities. *Journal of Public Health Policy* 1993;14(1):19–33.

Moore J. Non-profit advocates' latest tool: technology. *Chronicle of Philanthropy* 1994;6(24):1, 40–2.

Mullan F. Don Quixote, Machiavelli, and Robin Hood: public health practice, past and present. *American Journal of Public Health* 2000;90(5):702–6.

Murray DM, Hannan PJ, Jacobs DR, McGovern PJ, Schmid L, Baker WL, Gray C. Assessing intervention effects in the Minnesota Heart Health Program. *American Journal of Epidemiology* 1994;139(1):91–103.

Murray DM, Perry CL, Griffin G, Harty KC, Jacobs DR Jr, Schmid L, Daly K, Pallonen U. Results from a state-wide approach to adolescent tobacco use prevention. *Preventive Medicine* 1992;21(4):449–72.

National Cancer Institute. *ASSIST Program Guidelines for Tobacco-Free Communities*. Bethesda (MD): National Cancer Institute, Division of Cancer Prevention and Control, Cancer Control Science Program, Public Health Applications Research Branch, 1991.

National Cholesterol Education Program Expert Panel. Report of the National Cholesterol Education Program Expert Panel on detection, evaluation, and treatment of high blood cholesterol in adults. *Archives of Internal Medicine* 1988;148(1):36–69.

Nicholl J. *Tobacco Tax Campaign Guide*. Atlanta: American Cancer Society, 1996.

Nicholl J. Tobacco tax initiatives to prevent tobacco use: a study of eight statewide campaigns. *Cancer* 1998; 83(12 Suppl):2666–79.

Novotny TE, Siegel MB. California's tobacco control saga. *Health Affairs* 1996;15(1):58–72.

Nutbeam D. Achieving “best practice” in health promotion: improving the fit between research and practice. *Health Education Research* 1996;11(3):317–26.

Nutbeam D. Evaluating health promotion—progress, problems and solutions. *Health Promotion International* 1998;13(1):27–44.

Ockene JK, Lindsay EA, Hymowitz N, Giffen C, Purcell T, Pomrehn P, Pechacek T, for the COMMIT Research Group. Tobacco control activities of primary-care physicians in the Community Intervention Trial for Smoking Cessation. *Tobacco Control* 1997;6(Suppl 2):S49–S56.

Oregon Health Division. *Tobacco Prevention and Education Program Report—1999*. Portland (OR): Department of Human Resources, Oregon Health Division, 1999.

Orzechowski and Walker. *The Tax Burden on Tobacco: Historical Compilation* 1999. Arlington (VA): Orzechowski and Walker, 2000.

Pechacek TE, Lando HA, Nothwehr F, Lichtenstein E. Quit and Win: a community-wide approach to smoking cessation. *Tobacco Control* 1994;3(3):236–41.

- Pechmann C. Does antismoking advertising combat underage smoking? A review of past practices and research. In: Goldberg ME, Fishbein M, Middlestadt SE, editors. *Social Marketing: Theoretical and Practical Perspectives*. Mahwah (NJ): Lawrence Erlbaum Associates, 1997:189–216.
- Pechmann C, Reibling ET. Planning an effective anti-smoking mass media campaign targeting adolescents. *Journal of Public Health Management Practice* 2000; 6(3):80–94.
- Pechmann C, Shih CF. Smoking scenes in movies and antismoking advertisements before movies: effects on youth. *Journal of Marketing* 1999;63(3):1–13.
- Perry CL, Kelder SH, Murray DM, Klepp KI. Communitywide smoking prevention: long-term outcomes of the Minnesota Heart Health Program and the Class of 1989 Study. *American Journal of Public Health* 1992;82(9): 1210–6.
- Pertschuk M. *Smoke Signals: The Smoking Control Media Handbook*. Atlanta: American Cancer Society, 1987.
- Pertschuk M. SCARC: the Smoking Control Advocacy Resource Center. Developing a system for undergirding health policy advocacy. Paper presented at Progress Towards a Smokefree Society—National, State and Local Tobacco Control Initiatives, the American Public Health Association Annual Meeting; Nov 11, 1992; Washington (DC).
- Pertschuk M, Erickson A. *Smoke Fighting: A Smoking Control Movement Building Guide*. Atlanta: American Cancer Society, 1987.
- Pertschuk M, Wilbur P, O'Keefe AM. *Media Advocacy: Reframing Public Debate*. Washington: Benton Foundation, 1991.
- Pierce JP. International comparisons of trends in cigarette smoking prevalence. *American Journal of Public Health* 1989;79(2):152–7.
- Pierce JP. Mass media and tobacco control. In: Slama K, editor. *Tobacco and Health*. New York: Plenum Press, 1995:335–40.
- Pierce JP, Choi WS, Gilpin EA, Farkas AJ, Berry CC. Tobacco industry promotion of cigarettes and adolescent smoking. *Journal of the American Medical Association* 1998;279(7):511–5.
- Pierce JP, Dwyer T, Frape G, Chapman S, Chamberlain A, Burke N. Evaluation of the Sydney "Quit. For Life" anti-smoking campaign. Part 1. Achievement of intermediate goals. *Medical Journal of Australia* 1986;144(7):341–4.
- Pierce JP, Evans N, Farkas AJ, Cavin SW, Berry C, Kramer M, Kealey S, Rosbrook B, Choi W, Kaplan RM. *Tobacco Use in California: An Evaluation of the Tobacco Control Program, 1989–1993*. La Jolla (CA): University of California, San Diego, 1994.
- Pierce JP, Gilpin E, Burns DM, Whalen E, Rosbrook B, Shopland D, Johnson M. Does tobacco advertising target young people to start smoking? Evidence from California. *Journal of the American Medical Association* 1991;266(22):3154–8.
- Pierce JP, Gilpin EA, Emery SL, Farkas AJ, Zhu SH, Choi WS, Berry CC, Distefan JM, White MM, Soroko S, Navarro A. *Tobacco Control in California: Who's Winning the War? An Evaluation of the Tobacco Control Program, 1989–1996*. La Jolla (CA): University of California, San Diego, 1998a.
- Pierce JP, Gilpin EA, Emery SL, White MM, Rosbrook B, Berry CC. Has the California Tobacco Control Program reduced smoking? *Journal of the American Medical Association* 1998b;280(10):893–9.
- Pierce JP, Macaskill P, Hill D. Long-term effectiveness of mass media led antismoking campaigns in Australia. *American Journal of Public Health* 1990;80(5):565–9.
- Pollay RW, Siddarth S, Siegel M, Haddix A, Merritt RK, Giovino GA, Eriksen MP. The last straw? Cigarette advertising and realized market shares among youths and adults, 1979–1993. *Journal of Marketing* 1996;60:1–16.
- Puska P, Nissinen A, Salonen JT, Toumilehto [sic] J. Ten years of the North Karelia Project: results with community-based prevention of coronary heart disease. *Scandinavian Journal of Social Medicine* 1983a; 11(3):65–8.
- Puska P, Nissinen A, Tuomilehto J, Salonen JT, Koskela K, McAlister A, Kottke TE, Maccoby N, Farquhar JW. The community-based strategy to prevent coronary heart disease: conclusions from the ten years of the North Karelia project. *Annual Review of Public Health* 1985;6:147–93.

- Puska P, Salonen JT, Nissinen A, Tuomilehto J, Vartiainen E, Korhonen H, Tanskanen A, Rönqvist P, Koskela K, Huttunen J. Change in risk factors for coronary heart disease during 10 years of a community intervention programme (North Karelia project). *British Medical Journal* 1983b;287(6408):1840-4.
- Puska P, Tuomilehto J, Nissinen A, Vartiainen E. *The North Karelia Project: 20 Year Results and Experiences*. Helsinki (Finland): The National Public Health Institute (KTL), 1995.
- Puska P, Tuomilehto J, Salonen J, Neittaanmäki L, Maki J, Virtamo J, Nissinen A, Koskela K, Takalo T. Changes in coronary risk factors during comprehensive five-year community programme to control cardiovascular diseases (North Karelia project). *British Medical Journal* 1979;2(6199):1173-8.
- Riester T, Linton M. Designing an effective counter-advertising campaign—Arizona. *Cancer* 1998;83(12 Suppl):2746-51.
- Robert Wood Johnson Foundation. Nineteen states receive \$10 million to fight tobacco use [press release]. Princeton (NJ): Robert Wood Johnson Foundation, Aug 15, 1994.
- Robinson RG, Orleans CT, James DA, Sutton CD. *Pathways to Freedom: Winning the Fight Against Tobacco*. Philadelphia: Fox Chase Cancer Center, 1992. Reprinted with permission by the Centers for Disease Control and Prevention and the National Medical Association.
- Rogers EM. *Diffusion of Innovations*. 3rd ed. New York: Free Press, 1983.
- Rose G. Strategy of prevention: lessons from cardiovascular disease. *British Medical Journal Clinical Research Edition* 1981;282(6279):1847-51.
- Rose G, Heller RF, Pedoe HT, Christie DGS. Heart disease prevention project: a randomised controlled trial in industry. *British Medical Journal* 1980;280(6216):747-51.
- Ross M. *Tobacco Tax Campaigns: A Case Study of Two States*. Washington: Advocacy Institute, 1996.
- Rothman J. Three models of community organization practice, their mixing and phasing. In: Cox FM, Erlich JL, Rothman J, Tropman JE, editors. *Strategies of Community Organization*. 3rd ed. Itasca (IL): Peacock Publishers, 1979:25-45.
- Sallis JF, Flora JA, Fortmann SP, Taylor CB, Maccoby N. Mediated smoking cessation programs in the Stanford Five-City Project. *Addictive Behaviors* 1985;10(4):441-3.
- Salonen JT, Puska P, Kottke TE, Tuomilehto J. Changes in smoking, serum cholesterol and blood pressure levels during a community-based cardiovascular disease prevention program—the North Karelia Project. *American Journal of Epidemiology* 1981;114(1):81-94.
- Sanson-Fisher R, Redman S, Hancock L, Halpin S, Clarke P, Schofield M, Burton R, Hensley M, Gibberd R, Reid A, Walsh R, Girgis A, Burton L, McClintock A, Carter R, Donner A, Green S. Developing methodologies for evaluating community-wide health promotion. *Health Promotion International* 1996;11(3):227-36.
- Schmid TL, Pratt M, Howze E. Policy as intervention: environmental and policy approaches to the prevention of cardiovascular disease. *American Journal of Public Health* 1995;85(9):1207-11.
- Scientific Committee on Tobacco and Health. *Report of the Scientific Committee on Tobacco and Health*. London: The Stationery Office, 1998.
- Shea S, Basch CE. A review of five major community-based cardiovascular disease prevention programs. Part I: rationale, design, and theoretical framework. *American Journal of Health Promotion* 1990a;4(4):203-13.
- Shea S, Basch CE. A review of five major community-based cardiovascular disease prevention programs. Part II: intervention strategies, evaluation methods, and results. *American Journal of Health Promotion* 1990b;4(3):279-87.
- Shultz JM, Moen ME, Pechacek TF, Harty KC, Skubic MA, Gust SW, Dean AG. The Minnesota Plan for Non-smoking and Health: the legislative experience. *Journal of Public Health Policy* 1986;7(3):300-13.
- Siegel M. Mass media antismoking campaigns: a powerful tool for health promotion. *Annals of Internal Medicine* 1998;129(2):128-32.

- Siegel M, Biener L. The impact of an antismoking media campaign on progression to established smoking: results of a longitudinal youth study. *American Journal of Public Health* 2000;90(3):380-6.
- Siegel M, Mowery PD, Pechacek TF, Strauss WJ, Schooley MW, Merritt RK, Novotny TE, Giovino GA, Eriksen MP. Trends in adult cigarette smoking in California compared with the rest of the United States, 1978-1994. *American Journal of Public Health* 2000; 90(3):372-9.
- Specter M. Marketers target "virile female": R.J. Reynolds plans to introduce cigarette. *The Washington Post*, Feb 17, 1990;Sect A:1 (col 4).
- State of Arizona, Office of the Auditor General. *Performance Audit*. Department of Health Services' Tobacco Education and Prevention Program. Report to the Arizona Legislature. Phoenix (AZ): State of Arizona, Office of the Auditor General, 1999. Report No. 99-17.
- Steenkamp HJ, Jooste PL, Jordaan PCJ, Swanepoel ASP, Rossouw JE. Changes in smoking during a community-based cardiovascular disease intervention programme: the Coronary Risk Factor Study. *South African Medical Journal* 1991;79(5):250-3.
- Steinfeld JL. II: The public's responsibility: a bill of rights for the non-smoker. *Rhode Island Medical Journal* 1972;55(4):124-6, 138.
- Stockwell TF, Glantz SA. Tobacco use is increasing in popular films. *Tobacco Control* 1997;6(4):282-4.
- Substance Abuse and Mental Health Services Administration. *Synar Regulation: Report to Congress on FFY 1997 State Compliance*. Rockville (MD): US Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Prevention, 1998a. DHHS Publication No. (SMA) 97-3143.
- Substance Abuse and Mental Health Services Administration. *Synar Regulation: Report to Congress on FFY 1997 State Compliance: Summary*. Rockville (MD): US Department of Health and Human Services, Substance Abuse and Mental Health Services Administration, Center for Substance Abuse Prevention, 1998b. DHHS Publication No. (SMA) 98-3185.
- Susser M. The tribulations of trials—intervention in communities [editorial]. *American Journal of Public Health* 1995;85(2):156-8.
- Taylor HL, Blackburn H, Keys A, Parlin RW, Vasquez C, Puchner T. Coronary heart disease in seven countries. IV. Five-year follow-up of employees of selected U.S. railroad companies. *Circulation* 1970a;41 (4 Suppl): I20-I39.
- Taylor HL, Menotti A, Puddu V, Monti M, Keys A. Coronary heart disease in seven countries. XI. Five years of follow-up of railroad men in Italy. *Circulation* 1970b;41(4 Suppl):I113-I22.
- Taylor SM, Ross NA, Cummings KM, Glasgow RE, Goldsmith CH, Zanna MP, Corle DK. Community Intervention Trial for Smoking Cessation (COMMIT): changes in community attitudes toward cigarette smoking. *Health Education Research* 1998;13(1):109-22.
- Teenage Research Unlimited. *Counter-Tobacco Advertising Exploratory*. Summary Report, January-March 1999. Northbrook (IL): Teenage Research Unlimited, 1999.
- Teinowitz I. After the tobacco settlement; take the money & run (ads); the anti-smoking campaign's a many splendored thing, and that's the problem. *The Washington Post*, Dec 6, 1998; Outlook Sect: C1.
- Texas InterAgency Tobacco Task Force. *Legislative Plan*. Austin (TX): Texas Department of Health, 1998.
- Texas Department of Health. 1998 Texas Youth Tobacco Survey. Report 1: Current Tobacco Use; <<http://www.tdh.state.tx/us/otpc>>; accessed June 6, 2000.
- Thompson B, Corbett K, Bracht N, Pechacek T. Community mobilization for smoking cessation: lessons learned from COMMIT. *Health Promotion International* 1993;8(2):69-83.
- Tobacco Education and Research Oversight Committee. *Toward a Tobacco-Free California: Mastering the Challenges, 1995-1997*. Sacramento (CA): Tobacco Education and Research Oversight Committee, 1995.
- Tobacco Education Oversight Committee. *Toward a Tobacco-Free California: A Master Plan to Reduce Californians' Use of Tobacco*. Sacramento (CA): Tobacco Education Oversight Committee, 1991.